



Infants' Concentrated Drops 80 mg/0.8 mL	Children's Suspension Liquid 160 mg/5 mL	Children's Soft Chews Chewable Tablets 80 mg each	Junior Strength Chewable Tablets 160 mg each
--	--	---	--

Dose		Dropperful	Teaspoon (TSP)	Tablet	Tablet
WEIGHT	AGE				
6-11 lbs	0-3 mos	1/2 = (0.4mL)			
12-17 lbs	4-11 mos	1 = (0.8mL)	1/2 (TSP)		
18-23 lbs	12-23 mos	1-1/2 = (0.8 + 0.4mL)	3/4 (TSP)		
24-35 lbs	2-3 yrs	2 = (0.8 + 0.8mL)	1 (TSP)	2	
36-47 lbs	4-5 yrs		1-1/2 (TSP)	3	
48-59 lbs	6-8 yrs		2 (TSP)	4	2
60-71 lbs	9-10 yrs		2 1/2 (TSP)	5	2 1/2
72-95 lbs	11 yrs		3 (TSP)	6	3
96 lbs & over	12 yrs				4

Use only as directed.

NOTE: If possible, use weight to dose; otherwise use age. To arrive at the correct dose, weigh your child before giving TYLENOL®. All dosages may be repeated every 4 hours, but not more than 5 times daily.

A healthcare professional should be consulted for dosing for children under the age of two years.

WARNINGS:

- Children's TYLENOL® should not be taken for pain for more than 5 days or for fever for more than 3 days unless directed by a physician. If pain or fever persists or gets worse, if new symptoms occur, or if redness or swelling is present, a physician should be consulted because these could be signs of a serious condition.
- Do not exceed recommended dose. Taking more than the recommended dose (overdose) may not provide more relief and could cause serious health problems. Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a physician or poison control center immediately. Prompt medical attention is critical even if you do not notice any signs or symptoms.
- Do not use with any other product containing acetaminophen.
- Do not use Adult Extra Strength TYLENOL® products for children under 12 years of age.

Children's Motrin

IBUPROFEN

Ages 6 mos - 23 mos Infants' Motrin® Concentrated Drops 50 mg/1.25 mL	Ages 2 - 11 Children's Motrin® Suspension 100 mg/5 mL	Ages 2 - 11 Children's Motrin® Chewable Tablets 50 mg	Ages 6 - 11 Junior Strength Motrin® Chewable Tablets 100 mg	Ages 6 - 11 Junior Strength Motrin® Caplets 100 mg
--	--	--	--	---

Dose

Dropperful

Teaspoon (TSP)

Tablet

Tablet

Caplet

WEIGHT

AGE

Under 6 mos

Consult Your Child's Doctor

12-17 lbs

6-11 mos

1 - (1.25mL)

18-23 lbs

12-23 mos

1-1/2 - (1.875 mL)

24-35 lbs

2-3 yrs

1 (TSP)

2 tablets

36-47 lbs

4-5 yrs

1-1/2 (TSP)

3 tablets

48-59 lbs

6-8 yrs

2 (TSP)

4 tablets

2 tablets

2 caplets

60-71 lbs

9-10 yrs

2 1/2 (TSP)

5 tablets

2 1/2 tablets

2 1/2 caplets

72-95 lbs

11 yrs

3 (TSP)

6 tablets

3 tablets

3 caplets

Children's Motrin is available over-the-counter (OTC) for the temporary reduction of fever and relief of minor aches and pains due to colds, flu, sore throat, headaches and toothaches.

OTC Dosing: The recommended dose is 7.5 mg/kg every 6-8 hours. The recommended maximum daily dose is 30 mg/kg.

Keep all medication out of the reach of children.

Use only as directed.

References: 1. IMS NDTI Pediatricians recommendations, 12 months ending December 2000

2. IRI Data, YTD ending December 24, 2000.

McNEIL CONSUMER HEALTHCARE
DIVISION OF McNEIL - PPC, INC.
FORT WASHINGTON, PA 19034 USA